



ROBOTIC-ASSISTED LAPAROSCOPIC RADICAL PROSTATECTOMY (RALP)

Information about your procedure from
The British Association of Urological Surgeons (BAUS)

This leaflet contains evidence-based information about your proposed urological procedure. We have consulted specialist surgeons and patient groups during its preparation, so that it represents best practice in UK urology. You should use it in addition to any advice already given to you.

To view the online version of this leaflet, type the text below into your web browser:

[http://www.baus.org.uk/_userfiles/pages/files/Patients/Leaflets/Radical prostatectomy robot.pdf](http://www.baus.org.uk/_userfiles/pages/files/Patients/Leaflets/Radical%20prostatectomy%20robot.pdf)

Key Points

- This procedure aims to remove the prostate and seminal vesicles (sperm sacs) completely whilst trying to preserve the structures required to maintain urinary continence
- We use very small, robotic instruments that allow precise surgery through tiny “keyhole” incisions
- The instruments are totally under the control of the surgeon and the robot simply mimic the surgeon’s movements
- Erectile dysfunction (impotence), some leaking of urine and shortening of the penis can occur

What does this procedure involve?

This involves very precise removal of the whole prostate gland, seminal vesicles and, sometimes, the draining lymph glands, as well as tying off the vasa deferentia (sperm-carrying tubes). It is performed through several small puncture (“keyhole”) incisions in the lower abdomen (tummy) using robotic instruments.

Our aims in men with cancer confined to the prostate gland are:

- to remove the cancer,
- to achieve a clear margin away from the tumour
- to drop the PSA blood level below 0.1 ng per ml
- to reduce the need for any further treatment (e.g. radiotherapy or hormone treatment)
- to preserve your continence and,

- if possible and appropriate, to preserve the erection nerves to your penis.

Robotic surgery uses sophisticated mini-instruments which are totally under the control of the surgeon. The robot only mimics and assists the surgeon's movements; it does not do the operation. This technique is now widely used because of its high degree of surgical accuracy, and because recovery is much faster than it is for open surgery.

Your surgeon will try everything to preserve the muscle fibres and nerves that control continence. If you still leak some urine after a year (as 1 in 20 to 1 in 33 patients do), this can be corrected by another procedure such as an artificial urinary sphincter or a male sling.

The erection nerves are very close to the prostate, forming a cobweb of delicate strands over its surface. If your erections worked before the procedure, it is usually possible to preserve the nerves. This is called nerve-sparing prostatectomy. It can be very successful in maintaining your erections after the procedure although your erections may take some time to recover. We can only preserve these nerves if the cancer has not reached the layer where the nerve fibres lie.

Your team will explain how you can enjoy a healthy sex life after surgery, if the nerves do not recover or need to be removed.

What are the alternatives for organ-confined cancer?

- [**Active surveillance**](#) – no active treatment but careful monitoring of your PSA levels with repeated biopsies and further intervention only if there is definite evidence of cancer progression
- [**Open retropubic radical prostatectomy**](#) – performed through an incision in your lower abdomen
- [**Open perineal radical prostatectomy**](#) - performed through an incision just in front of your anus
- [**Standard laparoscopic \(keyhole\) radical prostatectomy**](#) – performed using the standard keyhole technique without robotic assistance
- [**External beam radiotherapy**](#) – giving an intensive course of external irradiation to your prostate gland
- [**Permanent seed brachytherapy**](#) – implanting radio-active seeds under ultrasound control into your prostate gland
- [**High intensity focused ultrasound \(HIFU\)**](#) – external beamed ultrasound; only available in a few specialist centres and, because we do not have long-term results, needs to be given as part of a clinical trial

- **Cryotherapy** – freezing & thawing the prostate with fine needles passed into the gland; only available in a few specialist centres and, because we do not have long-term results, needs to be given as part of a clinical trial

Deciding which treatment to have is not something you will do alone and may depend on the level of expertise available at your hospital. If you need further information, please contact your specialist nurse, surgical care practitioner or urologist.

You should be aware that there is a small chance (in 1 in 100 patients) that we might have to convert a laparoscopic (“keyhole”) procedure to open surgery. Because of this, we are not able to proceed with a robotic (keyhole) procedure if you are not willing to have open surgery under any circumstances.

What happens on the day of the procedure?

Your urologist (or a member of their team) will briefly review your history and medications, and will discuss the surgery again with you to confirm your consent.


You will be seen by an anaesthetist who will discuss the options of a general anaesthetic or spinal anaesthetic. The anaesthetist will also discuss pain relief after the procedure with you.

You may be given a pair of TED stockings to wear, and a heparin injection to thin your blood. These help to prevent blood clots from developing and from passing into your lungs. Your medical team will decide whether you need to continue these after you go home.

We normally give you a small enema before surgery to ensure that your bowel is clear.

Details of the procedure

- we will give you an injection of antibiotics into a vein before the procedure.
- once you are anaesthetised, the surgeon will make 5 or 6 keyhole cuts in your tummy (pictured) that allow the robotic instruments to be put in.
- these instruments allow the surgeon to free the prostate from the bladder and urethra (waterpipe) so it can be removed, whilst sparing



the muscles and nerves that control continence and trying to preserve your erection nerves.

- we then re-join your urethra to your bladder using absorbable stitches.
- we use local anaesthetic to numb the keyhole incisions so to minimise your discomfort when you wake up.
- all the keyhole incisions are closed with absorbable stitches.
- we will put a catheter in your bladder to drain the urine while the new join heals.
- the procedure usually takes two to three hours to perform.



After the procedure, you often get some bruising and swelling around the keyhole incisions together with some swelling or puffiness in your scrotum. You will often get some facial puffiness for a day or two (because you lie slightly “head down” during the surgery). You may also feel some shoulder pain and bloating until the bowel starts working again (normally after 24 hours).

Most patients go home after a day or two.

Are there any after-effects?

The possible after-effects and your risk of getting them are shown below. Some are self-limiting or reversible, but others are not. We have not listed very rare after-effects (occurring in less than 1 in 250 patients) individually. The impact of these after-effects can vary a lot from patient to patient; you should ask your surgeon's advice about the risks and their impact on you as an individual:

After-effect	Risk
No semen is produced during an orgasm, effectively making you infertile	 All patients
A high chance of erectile dysfunction (impotence) if a nerve-sparing operation is not possible or nerve damage is unavoidable, together with shortening of the penis	 Almost all patients
Urinary incontinence which may be temporary and require pads, or further surgery if lasts for more than one year	 Between 1 in 20 & 1 in 33 patients
Pathology tests that show cancer outside or at the margin of the prostate (positive margins) requiring observation & possible further treatment	 Between 1 in 10 & 1 in 50 patients
Further treatment with hormones, radiotherapy or chemotherapy may be needed at a later date if your PSA blood test still shows that cancer is present	 Between 1 in 10 & 1 in 50 patients
Leakage of urine from the new join between bladder & urethra, delaying discharge or needing longer catheter time	 Between 1 in 10 & 1 in 50 patients
Bleeding requiring transfusion or further surgery	 Between 1 in 10 & 1 in 50 patients
Pain, infection or hernia in any of the port incisions requiring further treatment	 Between 1 in 10 & 1 in 50 patients
	Between 1 in 10 &

Lymph fluid collection (if the pelvic lymph nodes were removed or biopsied during surgery)		1 in 50 patients
Anaesthetic or cardiovascular problems possibly requiring intensive care admission (including chest infection, pulmonary embolus, stroke, deep vein thrombosis, compartment syndrome, heart attack)		Between 1 in 50 & 1 in 250 patients (your anaesthetist can estimate your individual risk)
Need for conversion to open surgery due to operative difficulty or failure to progress		1 in 100 patients (1%)
Rectal injury or fistula requiring temporary colostomy to allow healing		Between 1 in 50 & 1 in 250 patients
Eye problems, or numbness & weakness due to nerve compression caused by your position during surgery		Between 1 in 50 & 1 in 250 patients

What is my risk of a hospital-acquired infection?

Your risk of getting an infection in hospital is approximately 8 in 100 (8%); this includes getting *MRSA* or a *Clostridium difficile* bowel infection. This figure is increased if you are in a “high-risk” group of patients such as patients who have had:

- long-term drainage tubes (e.g. catheters)
- bladder removal
- long hospital stays
- multiple hospital admissions

What can I expect when I get home?

- you will get some swelling and bruising of the incisions which may last several days
- it may be several days before you have your bowels open
- you may be discharged with a catheter in your bladder
- if you do have a catheter, we will show you how to manage it at home
- you will be given advice about your recovery at home
- you will be given a copy of your discharge summary and a copy will also be sent to your GP
- any antibiotics or other tablets you may need will be arranged & dispensed from the hospital pharmacy

- a follow-up appointment will be made for you to have your dressings and your catheter removed
- once your catheter has been removed, you should start doing [pelvic floor exercises](#)
- do not worry if you leak some urine when your catheter comes out; almost everyone has a period of bladder recovery when they will need to wear protective pads
- we will discuss the microscopic analysis of your prostate in a multi-disciplinary team (MDT) meeting
- we will arrange for you to have your first PSA check six to eight weeks after the procedure

General information about surgical procedures

Before your procedure

Please tell a member of the medical team if you have:

- an implanted foreign body (stent, joint replacement, pacemaker, heart valve, blood vessel graft)
- a regular prescription for a blood thinning agent (warfarin, aspirin, clopidogrel, rivaroxaban or dabigatran)
- a present or previous MRSA infection
- a high risk of variant-CJD (e.g. if you have had a corneal transplant, a neurosurgical dural transplant or human growth hormone treatment)

Questions you may wish to ask

If you wish to learn more about what will happen, you can find a list of suggested questions called "[Having An Operation](#)" on the website of the Royal College of Surgeons of England. You may also wish to ask your surgeon for his/her personal results and experience with this procedure.

BAUS runs a national audit and collects data from all urologists undertaking this surgery. There are two reasons for this. First, surgeons are required by the Department of Health to look at how well the surgery is being done under their care and, second, to look at national trends for the procedure.

Some basic patient data (e.g. name, NHS number and date of birth) are entered and securely stored. This is required so that members of the clinical team providing your care can go back to the record and add follow-up data such as length of stay or post-operative complications. This helps your surgeon to understand the various outcomes of the procedure.

Although BAUS staff can download the surgical data for analysis, they **cannot** access any patient identifiable data. This information is used to

generate reports on individual surgeons and units; these are available for the public to view in [Surgical Outcomes Audit](#) section of the BAUS website.

Before you go home

You should be told how the procedure went and you should:

- make sure you understand what has been done
- ask the surgeon if everything went as planned
- let the staff know if you have any discomfort
- ask what you can (and cannot) do at home
- make sure you know what happens next
- ask when you can return to normal activities

You will be given advice about what to look out for when you get home. Your surgeon or nurse will also give you details of who to contact, and how to contact them, in the event of problems.

Smoking and surgery

Ideally, we would prefer you to stop smoking before any procedure. Smoking can worsen some urological conditions and makes complications more likely after surgery. For advice on stopping, you can:

- contact your GP
- access your local [NHS Smoking Help Online](#)
- ring the free NHS Smoking Helpline on **0800 169 0 169**

Driving after surgery

It is your responsibility to make sure you are fit to drive after any surgical procedure. You only need to [contact the DVLA](#) if your ability to drive is likely to be affected for more than three months. If it is, you should check with your insurance company before driving again.

What should I do with this information?

Thank you for taking the trouble to read this information. Please let your urologist (or specialist nurse) know if you would like to have a copy for your own records. If you wish, they can also arrange for a copy to be kept in your hospital notes.

What sources were used to prepare this leaflet?

This leaflet uses information from consensus panels and other evidence-based sources including:

- the [Department of Health \(England\)](#)
- the [Cochrane Collaboration](#)
- the [National Institute for Health and Care Excellence \(NICE\)](#)

It also follows guidelines from:

- the [Royal National Institute for Blind People \(RNIB\)](#)
- the [Patient Information Forum](#)
- the [Plain English Campaign](#)

Disclaimer

We have made every effort to give accurate information but there may still be errors or omissions in this leaflet. BAUS cannot accept responsibility for any loss from action taken (or not taken) as a result of this information.

PLEASE NOTE

The staff at BAUS are not medically trained, and are unable to answer questions about the information provided in this leaflet. If you do have any questions, you should contact your urologist, specialist nurse or GP.